On measures of nonplanarity of cubic graphs

Leonid Plachta

Abstract. We study two measures of nonplanarity of cubic graphs G , the genus $\gamma(G)$, and the edge deletion number $ed(G)$. For cubic graphs of small orders these parameters are compared with another measure of nonplanarity, the rectilinear crossing number $\overline{cr}(G)$. We introduce operations of connected sum, specified for cubic graphs G , and show that under certain conditions the parameters $\gamma(G)$ and $ed(G)$ are additive (subadditive) with respect to them.

The minimal genus graphs (i.e. the cubic graphs of minimum order with given value of genus γ) and the minimal edge deletion graphs (i.e. cubic graphs of minimum order with given value of edge deletion number ed) are introduced and studied. We provide upper bounds for the order of minimal genus and minimal edge deletion graphs.

Анотація. З відомої теореми Куратовського випливає, що кубічний граф є непланарним тоді і тільки тоді, коли він не містить підграфів, гомеоморфних $K_{3,3}$. Для непланарних графів існує декілька характеристик графа, які визначають міру його непланарності. Для заданого 3-зв'язного кубічного графу G позначимо через $ed(G)$ найменше число ребер в G , після викидання яких дістанемо планарний підграф, а через $g(G)$ (орієнтовний) рід графа G . Крім того, нехай $\overline{cr}(G)$ позначає мінімальне число внутрішніх перетинів ребер графа G серед усіх прямолінійних імерсій графа в площині. Кубічний граф G називається k -мінімальним відносно параметра $ed(G)$ (відповідно, параметрів $\overline{cr}(G)$, $g(G)$), якщо $ed(G) = k$ (відповідно, $\overline{cr}(G) = k$, $g(G) = k$) і порядок графа G є мінімальним серед усіх 3-зв'язних кубічних графів з даною властивістю.

В роботі досліджуються k -мінімальні відносно параметрів $ed(G)$ і $g(G)$ 3-зв'язних кубічних графи. Описані операції на 2-зв'язних і 3-зв'язних кубічних графах (зв'язна сума, подвійна зв'язна сума), які мають властивість адитивності (субадитивності) відносно параметрів $ed(G)$ і $g(G)$.

The research partially supported by the Polish Ministry of Science and Higher Education
2010 Mathematics Subject Classification: 05C10, 05C15

Keywords: nonplanar graphs, cubic graphs, genus, edge deletion number, connected sum of graphs, minimal graphs

Обчислені характеристики $ed(G)$ і $g(G)$ k -мінімальних відносно параметра $\overline{cr}(G)$ кубічних графів для малих чисел k . Дається порівняння характеристик $ed(G)$, $g(G)$ і $\overline{cr}(G)$ для зв'язних кубічних графів G .

Запропонований конструктивний метод, який дозволяє отримувати 2-зв'язні і циклічно 4-зв'язні кубічні графи G з як завгодно великими характеристиками $ed(G)$ і $g(G)$. Даний метод базується на адитивних властивостях операцій “зв'язна сума” і “подвійна зв'язна сума” на кубічних графах. Отримані також верхні оцінки порядку мінімальних відносно характеристик $ed(G)$ і $g(G)$ графів G , в класі 2-зв'язних і 3-зв'язних кубічних графів G . Сформульована також відкрита проблема, яка стосується нижньої оцінки для порядку k -мінімальних відносно параметрів $ed(G)$ і $g(G)$ кубічних графів.

1. INTRODUCTION

We consider finite graphs $G = (V, E)$ without loops and multiple edges. The number of vertices of G is called the *order* of G and denoted by $|G|$. The number of edges $|E|$ of the graph G is called its *size* and denoted by $||G||$. The Kuratowski theorem states that a graph G is planar if and only if it does not contain subgraphs homeomorphic to K_5 and $K_{3,3}$. A regular graph of valence 3 is called *cubic*. For cubic graphs, the only forbidden graphs are those which are not homeomorphic to $K_{3,3}$. There are different measures of nonplanarity of a graph. Let us recall their definitions.

For a given connected graph G denote by $\gamma(G)$ the (*orientable*) *genus* of G i.e. the minimal genus of an orientable closed connected surface M such that G has an embedding in M . Note that each such embedding is 2-cell. The problem of deciding whether a cubic graph G has the genus $\gamma(G) \leq m$ is known to be NP-complete, [21]. There are some upper and lower bounds of $\gamma(G)$ for different classes of graphs G , [19]. For cubic graphs G , the precise values of the parameter $\gamma(G)$ are known only for special classes of them (for example, for some snarks, etc., see [15, 19]).

Another well known measure of nonplanarity of a graph G is the *crossing number* $cr(G)$ (the *rectilinear crossing number* $\overline{cr}(G)$). This is the minimal number of proper double crossings of edges among all immersions of G in the plane (the minimal number of proper double crossings of edges among all rectilinear immersions of G in the plane, respectively). The computation of the crossing number of a graph is also an NP-complete problem, [7]. Note that, in general, $cr(G)$ and $\overline{cr}(G)$ are distinct numbers, [2]. There are estimations of the parameters $cr(G)$ and $\overline{cr}(G)$ for complete graphs, complete bipartite graphs, and other special classes of graphs (see, for example [10, 20]). The precise values of $cr(G)$ and $\overline{cr}(G)$ are known only for particular nonplanar graphs (for example, for small complete and complete bipartite graphs, [17, 20]).

For a given graph G , denote by $ed(G)$ the minimal number of edges in G such that after their deletion the resulting graph becomes planar. The parameter $ed(G)$ is called *the edge deletion number of G* and the corresponding problem of finding the minimal set of edges to be deleted in a graph G is known as MINED. Even for cubic graphs, the problem MINED is known to be NP-complete, [5]. Algorithms of computing $ed(G)$, in particular, for cubic graphs, are described in [3, 5, 4].

Comparing with the parameters $\gamma(G)$ and $cr(G)$, there are much more fewer results concerning evaluation of the number $ed(G)$. Pegg jr and Exoo [12] introduced the notion of a minimal crossing graph. For a given natural number k a cubic graph G is called *minimal k -crossing graph* (in original, k -crossing graph [12]) if G has a minimal order among all cubic graph H with $\overline{cr}(H) = k$. By this analogy, we introduce minimal k -genus and minimal k -edge deletion graphs. Denote by $\beta(G)$ the *cyclomatic* number of the connected graph G , $\kappa(G)$ the *vertex connectivity* and $\lambda(G)$ the *edge connectivity* of G . For graphs G of maximal degree at most three the number $\lambda(G)$ and $\kappa(G)$ coincide, [6]. By this reason, for cubic graphs G we will abbreviate the terminology and use the term “the connectivity of G ”.

We say that a connected graph $G = (V, E)$ is *cyclically k -edge connected* if no set of fewer than k edges is cycle-separating in G . The *edge cyclic connectivity* $\zeta(G)$ of the cubic graph G is the largest integer $k \leq \beta(G)$ for which G is cyclically k -edge connected. For any cubic connected graph G we have obviously $\kappa(G) = \lambda(G) \leq \zeta(G)$. Note that $\zeta(G)$ is equal to $\beta(G)$ if and only if G does not have any cycle-separating edge cut. Moreover for cubic graphs G with $\zeta(G) \leq 3$ the values of vertex connectivity, edge connectivity and cyclic k -edge connectivity coincide, [16]. As an example, for the Petersen graph P we have $\kappa(P) = \lambda(P) = 3$ but $\zeta(P) = 5$.

We say that a connected cubic graph G is *cyclically k -vertex connected* if it contains no cycle-separating vertex cut with fewer than k vertices. For exception of few graphs (which are $K_4, K_{3,3}$ and the multigraph Θ_2), the notions of cyclically k -vertex connected graph and cyclically k -edge connected graph coincide, [16, Proposition 3].

In Section 2, we evaluate genus and edge deletion number of minimal k -crossing graphs for small numbers k . These auxiliary results are used in Section 3.

Battle et al. [1] have shown that the genus of any connected graph is equal to the sum of blocks with respect to its block decomposition. This is perhaps the first known result on additivity of the (orientable) genus of graph. The operation of the vertex amalgamation applied to 2-connected cubic graphs gives a separable graph which contains a vertex of degree 4.

Another operation is the edge amalgamation of graphs G_1 and G_2 , [13]. Miller, [13], introduced the generalized genus of a graph and showed that it is additive with respect to the edge amalgamation of two graphs. The operation of edge amalgamation does not preserve the class of cubic graphs. In [8], Gross also studied bar-amalgamation of graphs. All these operations, when applied to cubic graphs, produce the graphs which are outside the given class. Moreover they are not compatible with such properties of graphs (including cubic graphs) as the vertex connectivity and the edge connectivity. Therefore they cannot serve as a good tool for the construction of graphs with big numbers of the parameters ed and γ inside the class of cubic graphs with the given connectivity.

In Section 3, we introduce two operations of connected sum which are suitable for cubic graphs. We study additivity properties of genus and edge deletion number with respect to these operations. The first operation, when applied to two 2-connected cubic graphs, results in a 2-connected cubic graph. Similarly, the second operation preserves, in general, the class of 3-connected (or even cyclically 4-edge connected) cubic graphs. Additivity properties of cubic graphs are provided by Theorems 3.1, 3.2 and 3.3 (subject to the parameter γ), and by Theorems 3.4 and 3.5 (subject to the parameter ed). By using these properties, we provide upper bounds for the order of a 2-connected and 3-connected cubic graphs G , which are minimal with respect to these parameters (Corollaries 3.1-3.4). These are the main results of the paper.

In [18] we use more subtle arguments for obtaining upper bounds of the order of minimal cubic graphs G with prescribed value of the parameter γ . The additivity properties of the parameter γ given by Theorems 3.1, 3.2 and 3.3 are also essential in this relation.

2. MEASURES OF NONPLANARITY OF CUBIC GRAPHS: SMALL ORDERS

We start by considering the parameters $cr(G)$ and $\overline{cr}(G)$ for small cubic graphs G and compare these numbers with the parameters $\gamma(G)$ and $ed(G)$. Denote by $g(G)$ the girth of the graph G . In our study of the (orientable) genus of cubic graphs we shall use the notion of the rotation system on a graph.

A *rotation system* on a graph $G = (V, E)$ is a family $\Pi = \{\pi_v : v \in V\}$, where π_v is a cyclic permutation of the edges incident with v . With any 2-cell embedding φ of a graph G into an oriented closed surface S it is associated a rotation system Π on G . The pair (G, Π) is called a *rotation graph*. Moreover for a given rotation graph (G, Π) one can construct a system R of (oriented) circuits on G in such a way that each edge e of G is contained twice in the circuits, but with opposite orientations. The circuits

c from R can be thought of as oriented boundaries of 2-dimensional discs D_c . Gluing together the family of discs $D_c, c \in R$, along their oriented boundaries, we shall obtain an oriented closed surface S and this provides a natural 2-cell embedding ψ of G into S . The circuits c from R are called the *facial cycles* of the embedding ψ . The correspondence between the 2-cell embeddings of a graph G into oriented surfaces and rotation systems on G is one-to-one, in a usual sense. For more details see [9], [14].

It is easy to see that we have the following inequalities:

$$\gamma(G) \leq ed(G) \leq cr(G) \leq \overline{cr}(G).$$

It can be shown that for cubic graphs the difference between any two of the parameters $\gamma(G), ed(G), cr(G)$ of G can be arbitrarily large. This can be proven, for example, by using results of Sections 2 and 3. Moreover, there exist graphs G for which the number $cr(G)$ is less than $\overline{cr}(G)$ (more precisely, $cr(G) = 4$ and $\overline{cr}(G) = m$ for any $m > 4$, [2]).

We shall say that a cubic graph G is *minimal l -genus graph* if $\gamma(G) = l$ and it is of minimum order among all 2-connected cubic graphs with this property. Similarly, for a given nonnegative integer l , a cubic graph G is *minimal l -edge deletion graph*, if $ed(G) = l$ and G is of minimum order among all 2-connected cubic graphs with this property.

In this section, we evaluate or estimate the order of minimal graphs with respect to parameters γ and ed for small numbers l . First count all minimal l -crossing graphs G for small values l . Minimal l -crossing graphs have been described up to value $l \leq 8$ in [12]. Note that for $l = 9$ it is unknown any minimal crossing graph G . At present, for $l \geq 10$, there are known only hypothetically minimal l -crossing graphs. Using minimal l -crossing graphs, we will find some minimal cubic graphs with respect to parameters ed and γ . For cubic graphs of small order we use the notations as in [12].

In the following, we will work in the piece wise linear category PL, [11]. Therefore surfaces are 2-dimensional PL-manifolds, graphs are 1-dimensional polyhedra, the maps (embeddings) of graphs are PL-maps (PL-embeddings) and the images of graphs under such maps are subpolyhedra of PL-manifolds (surfaces). For more detailed information about the category PL see also [14].

1. For $l = 1$ there is a unique minimal crossing graph, the graph $K_{3,3}$. We have obviously

$$ed(K_{3,3}) = \overline{cr}(K_{3,3}) = cr(K_{3,3}) = \gamma(K_{3,3}) = 1.$$

2. For $l = 2$ there are two minimal crossing graphs. These are the Petersen graph P (see Figure 2.1b) and the graph $CNG2B$ (see Figure 2.1a). We have obviously

$$ed(P) = 2, \quad \gamma(P) = 1, \quad ed(CNG2B) = \gamma(CNG2B) = 1.$$

FIGURE 2.1. The minimal 2-crossing graphs

We shall say that a system F of (oriented) circuits in a graph G is *admissible* if it determines a rotation system or can be completed to a rotation system on G by adding some additional circuits in G . In the latter case, we will say that F is *incomplete*. Note that for a connected cubic graph G of order 12, any incomplete admissible system F on G consisting of 4 circuits can be completed to a rotation system R that induces embedding of G in torus or the sphere S^2 .

Lemma 2.1. *For any connected cubic graph G of order 12 we have*

$$\gamma(G) \leq 1.$$

Proof. Let G be a connected cubic graph of order 12. We know from [12] that if $|G| \leq 12$, then $\overline{cr}(G) \leq 2$. If $\kappa(G) = 1$, the assertion follows (by using for example results from [1]). If $\kappa(G) = 2$, then after removal two nonincident edges from G we shall obtain two planar subcubic graphs, G_1 and G_2 . The only case, when one can worry about, is that G_1 and G_2 are isomorphic to $K_{3,3} - e$ where e is an edge of $K_{3,3}$. But in this case, G is isomorphic to a connected sum of two copies of $K_{3,3}$ and has genus 1 (see Section 3). Therefore we may assume that G is 3-connected. If $\overline{cr}(G) = 1$ we have obviously $\gamma(G) = 1$. If $\overline{cr}(G) = 2$ and the equality reaches via a straight line drawing G in the plane, in which one edge intersects two another edges, the assertion also easily follows.

Assume that there is an immersion of G in the oriented plane P in which we have crossings of two pairs of different edges: e_1 and e_2 , and f_1 and f_2 . Deleting e_1, e_2, f_1 and f_2 from G , we get a subcubic multigraph H which has a natural embedding φ in P . We may assume without loss of generality that H is connected, otherwise one can use a flip and redraw G in the plane with a fewer number of crossings. Denote by Π the rotation system on H associated with the embedding φ . Now consider all possible configurations of the induced plane embedding of the (multi)graph H and the positions of the deleted edges with respect to it.

a) There is an inner face r of the embedding φ which contains two pairs of crossing edges, e_1 and e_2 , and f_1 and f_2 . We have three types of configurations describing positions of these edges inside a *regular* face r . In

any case, we are able to define an admissible system of circuits in G which, after a suitable completion, generates an embedding of G into torus.

In the first case (see Figure 2.2), we have inside r two internal vertices of G and two crossings of different pairs of edges. The position of external two edges of H and incident vertices is irrelevant and so, is not indicated here. One can replace the facial circuit dr of the rotation system Π with three new circuits c_1, c_2, c_3 as indicated in Figure 2.2. The system of circuits $F = \{c_1, c_2, c_3\}$ in G is admissible and contain each “internal” edge of the immersed graph G twice, but with opposite orientations. The orientation of edges positioned on dr coincide with the one of the cycle dr . We can complete F to a rotation system on G by adding three outer circuits of Π , u_1, u_2 and u_3 . The system of circuits $\mathcal{F} = \{c_1, c_2, c_3, u_1, u_2, u_3\}$ defines a rotation system T on G of genus 1.

FIGURE 2.2.

In the second case, we have inside r one internal vertex of G and two crossings of edges, as shown in Figure 2.3a). The edges of G positioned inside r are called internal while the edges in the exterior of r are called external (subject to the given immersion of G in P). Let E' be the set of external edges of G . The vertices of G incident to external edges are called external. Denote by H_1 the graph $G - E'$. Then H_1 is immersed via φ in the closure of the face r . Consider in G the system of circuits

$$F_1 = \{c_1, c_2, c_3, c_4, c_5, c_6\}$$

indicated in Figure 2.3b). F_1 defines an embedding of H_1 into a sphere S^2 . We have two possibilities:

- a) G is obtained from H_1 by gluing two nonincident edges e_1 and e_2 to H_1 ;
- b) G is obtained from H_1 by gluing the graph $K_{1,3}$ along three vertices of degree 1.

Consider the subcase a). Suppose that the end vertices of e_1 (or e_2) are lying on the same circuit c_i . One can replace c_i in the system F_1 with two new circuits, c'_i and c''_i , as shown in Figure 2.3c). Denote by F_2 the new

FIGURE 2.3.

family consisting of seven circuits. Obviously F_2 is a rotation system on the graph $H_1 \cup e_1$ ($H_1 \cup e_2$, respectively). The remaining edge e_k from $\{e_1, e_2\}$ joins the vertices of different circuits of F_2 , say d_i and d_j . This results in a single circuit d in G containing both d_1 and d_2 and the edge e_k , which is involved twice and with opposite directions. Finally we get a rotation system R on G consisting of six circuits.

Now suppose that both e_1 and e_2 joins the vertices of the same circuits, c_i and c_j of F_1 . The vertices of e_1 and e_2 divide c_i into two directed paths, l_1 and l_2 , and c_j into two directed paths, m_1 and m_2 . Combining the paths l_1, l_2, m_1 and m_2 with the oriented edges e_1 and e_2 , we get two circuits in G , d_i and d_j . Each edge $e_i, i = 1, 2$, is contained both in d_i and d_j , but with opposite orientation. Replacing the pair of circuits c_i and c_j with the pair d_i and d_j , we obtain an admissible system F in G , consisting of six circuits. Therefore F defines in G a rotation system of genus one.

In the remaining cases, we can take three circles c'_1, c'_2 and c'_3 of F_1 , endow them with a suitable orientation, and add to them a new circuit c of G , which contains e_1 or e_2 . This results in an admissible system F' on G consisting of 4 circuits. The choice of c'_1, c'_2 and c'_3 depends on a position of vertices of e_1 or e_2 on dr (see, for example, Figure 2.3d). Completing F' by two new circuits, we get a rotation system R on G of genus one.

The subcase b) is handled in the same way as subcase a). We omit here the details.

The third case of configuration for H_1 is indicated in Figure 2.4a). We also have two possibilities:

- G is obtained from H_1 by gluing up two nonincident edges e_1 and e_2 ;
- G is obtained from H_1 by gluing up the graph $K_{1,3}$ along three vertices of degree 1.

There is no essential difference between the subcases a) and b). Let us consider the subcase a). We can choose two circuits c_1 and c_2 in H_1 and add to them two circuits d_1 and d_2 in G such that d_1 contains the edge e_1

and d_2 contains the edge e_2 . As a result, we get an incomplete admissible system F' of circuits in G . The choice of circuits c_1 and c_2 depends on the position of vertices of the edges e_1 and e_2 on dr , see Figure 2.4b) and c). Then F' can be completed to an admissible system F in G , consisting of six circuits. An exceptional case of the pairs of crossing edges e_1, e_2 and f_1, f_2

FIGURE 2.4.

inside a *nonregular* face r is shown in Figure 2.5. The following family of circuits in G determines a rotation system R of genus one:

$$\begin{aligned} c_1 &= (v_1, v_2, v_3, v_4), & c_2 &= (v_4, v_3, v_5, v_6, u_6), \\ c_3 &= (v_6, u_2, u_3, u_5, u_6), & c_4 &= (u_2, u_1, u_4, u_3), \\ c_5 &= (u_1, u_2, v_6, v_5, v_1, v_4, u_6, u_5), & c_6 &= (u_1, u_5, u_3, u_4, v_2, v_1, v_5, v_3, v_2, u_4). \end{aligned}$$

FIGURE 2.5.

b) There are two faces r_1 and r_2 of the embedding φ such that r_1 contains the crossing of e_1 and e_2 , and r_2 contains the crossing of f_1 and f_2 .

If r_1 and r_2 are disjoint, the existence of a rotation system Π' on G with 6 circuits is obvious. If r_1 and r_2 have a unique edge in common, we have a configuration shown in Figure 2.6. There is a rotation system R on G with 6 facial circuits. We indicate here only a noncomplete admissible system consisting of four circuits, $F = \{c_1, c_2, c_3, c_4\}$. The circuits are the following:

$$c_1 = (u_2, u_3, u_4, u_1), \quad c_2 = (u_1, u_4, u_5, u_{11}, u_{12}),$$

$$c_3 = (u_{11}, u_5, u_6, u_9, u_{10}), \quad c_4 = (u_9, u_6, u_7, u_8).$$

FIGURE 2.6.

Assume now that r_1 and r_2 have two edges in common, see Figure 2.7. In this case, we indicate a rotation system R on G with the following six circuits:

$$\begin{aligned} c_1 &= (u_2, u_1, v_2, v_1), & c_2 &= (v_5, v_4, u_5, u_4), \\ c_3 &= (u_1, u_2, u_3, u_4, u_5, u_6), & c_4 &= (v_1, v_2, v_3, v_4, v_5, v_6), \\ c_5 &= (u_2, v_1, v_6, u_6, u_5, v_4, v_3, u_3), & c_6 &= (u_1, u_6, v_6, v_5, u_4, u_3, v_2, v_1). \end{aligned}$$

FIGURE 2.7.

c) It can occur that H is a multigraph with three loops and the pairs of crossing edges of G are situated in the outer face p of the embedding φ . We depict in Figure 2.8 such a configuration. In this case, we indicate the following noncomplete admissible system of circuits in G :

$$\begin{aligned} c_1 &= (u_{10}, u_9, u_{12}, u_{11}), & c_2 &= (u_{11}, u_{12}, u_3, u_4), \\ c_3 &= (u_4, u_3, u_2, u_1), & c_4 &= (u_5, u_6, u_7). \end{aligned}$$

Note that the case when one pair of crossing edges of G is inside p and the other one is inside a region bounded by a loop of H is not admissible by the assumption that the graph G is 2-connected. Lemma 2.1 is completed. $\square$

FIGURE 2.8.

3. For $l = 3$ there are eight minimal crossing graphs. We count them according to [12]: *CNG 3A*, *CNG 3B*, *CNG 3D*, *CNG 3E*, *CNG 3F*, *CNG 3H*, *GP(7, 2)*, the Heawood graph *H* (see Figure 2.9).

FIGURE 2.9. The minimal 3-crossing graphs

By direct computation, we have

$$\begin{aligned} ed(CNG\ 3A) &= ed(CNG\ 3B) = ed(CNG\ 3E) \\ &= ed(CNG\ 3F) = ed(CNG\ 3H) = 2. \end{aligned}$$

In Figure 2.9, for each of these graphs we indicate by bold line the two edges after removal of which we obtain a planar subgraph. In Figure 2.10a), we indicate 3-crossing drawing of the graph *GP(7, 2)*. By bold lines there are indicated two edges in *GP(7, 2)* after removal of which we obtain a planar subgraph. Similarly, in Figure 2.10b) we indicate 3-crossing drawing of the graph *CNG 3D*. Removing two bold edges from it also leads to a planar subgraph.

FIGURE 2.10.

Lemma 2.2. *For the Heawood graph H we have $ed(H) = 3$.*

Proof. It is well known that H is symmetric graph. Remove any edge e from H . The resulting graph U can be represented as a sum of two subgraphs H_1 and H_2 and the edge f , see Figure 2.11. The subgraphs H_1 and H_2 have in common a path l of length 5. Denote by $E(l)$ the set of edges of the path l .

FIGURE 2.11.

Removing the edge f from U , we obtain obviously a non planar graph. Let h be any edge of the graph $H_1 \setminus E(l)$. There is a path $p_1 \subset H_1 \setminus \{E(l) \cup h\}$ such that $H_2 \cup p_1$ contains a graph homeomorphic to $K_{3,3}$. Similarly, for any edge g of the graph $H_2 \setminus E(l)$ there is a path $p_2 \subset H_2 \setminus \{E(l) \cup g\}$ such that $H_1 \cup p_2$ contains a graph homeomorphic to $K_{3,3}$. Therefore removing any edge k from $U \setminus E(l)$ results in a non planar subgraph. On the other hand, it is not difficult to check that removing any edge $e_i \in E(l)$ from U also leads to a graph which contains a subgraph homeomorphic to $K_{3,3}$. As an example consider the subgraph $U \setminus e_5$ shown in Figure 2.12. The subgraph homeomorphic to $K_{3,3}$ is depicted here by bold line. Therefore $ed(H) \geq 3$. Since $cr(H) = 3$ it follows that $ed(H) = 3$. $\square$

It is known that the Heawood graph H is toroidal, [19]. The Heawood graph is also cyclically 4-edge connected. To show this we first note that H is symmetric. Remove an edge e from H as in the proof of Lemma 2.2.

FIGURE 2.12.

Ignoring two vertices of degree 2 in the resulting graph U' we shall obtain a cubic graph U of degree 12, which is homeomorphic to U' . Evidently, U is 3-connected. It follows that U is cyclically 3-edge connected, so it H is cyclically 4-edge connected. The last fact will be used in Section 3.

Note also that $\gamma(CNG\ 3A) = 2$. The proof of this fact will be given in Section 3. It follows that $CNG\ 3A$ is a minimal 2-genus graph. It is not difficult to check that $\zeta(CNG\ 3A) = 3$. By direct computation, the remaining seven 3-crossing graphs have genus equal to one. We omit here the details of this computation.

4. For $l = 4$ there are two minimal crossing graphs: 8-crossed prism graph Pr_8 , see Figure 2.13a), and the Möbius-Kantor graph MK , see Figure 2.13b). By direct computation we have $ed(MK) = 3$ and $ed(Pr_8) = 2$. Moreover it is known that the Möbius-Kantor graph MK is toroidal, [12]. It is not difficult to show that the graph Pr_8 is also toroidal.

FIGURE 2.13. Graphs Pr_8 and MK

3. ADDITIVITY OF PARAMETERS γ AND ed AND MINIMAL CUBIC GRAPHS

In this section we introduce two operations on graphs and establish some additivity properties of parameters ed and γ with respect to them, in the case of cubic graphs. The first operation is the connected sum of graphs and the second one is the double (crossed) connected sum of them. We also provide some upper bounds for the order of minimal edge deletion and

minimal genus graphs within the classes of 2-connected and 3-connected cubic graphs.

Let G_1 and G_2 be 2-connected cubic graphs with distinguished edges e in G_1 and f in G_2 . Let u_1, u_2 be the vertices of e and v_1, v_2 the vertices of f , respectively. Remove from G_1 the edge e , and from G_2 the edge f . Take the disjoint sum G of resulting graphs, $G = (G_1 - e) \sqcup (G_2 - f)$, and joint in G the pairs of vertices: u_1 with v_1 , and u_2 with v_2 , respectively. Denote the resulting graph by $G_1 \star G_2$. We shall say that $G_1 \star G_2$ is *the connected sum* of the graphs G_1 and G_2 with respect to the pair of edges e and f . Note $G_1 \star G_2$ is also 2-connected cubic graph.

Let G_1 and G_2 be any two 3-connected graphs. Take in G_1 a pair of nonincident edges (e_1, e_2) , and in G_2 a pair of nonincident edges (f_1, f_2) . Denote the vertices of e_1 by u_1, u_2 , and the vertices of e_2 by v_1, v_2 , respectively. Similarly, let s_1, s_2 be the vertices of f_1 , and t_1, t_2 the vertices of f_2 . Delete in G_1 the edges e_1 and e_2 , and in G_2 the edges f_1 and f_2 . Then take a disjoint sum $G = (G_1 - e_1 - e_2) \sqcup (G_2 - f_1 - f_2)$ of two graphs and joint in G the following pairs of vertices: u_1 and s_1 , u_2 and s_2 , v_1 and t_1 , and v_2 and t_2 , respectively. Denote the resulting 2-connected graph $G_1 \star G_2$ and call it *a double connected sum* of G_1 and G_2 . The four edges joining the graphs $G_1 - e_1 - e_2$ and $G_2 - f_1 - f_2$ are called the bridge edges of the graph $G_1 \star G_2$ and are denoted h_1, h_2, h_3 and h_4 , see Figure 3.1.

FIGURE 3.1. A double connected sum of graphs G_1 and G_2

If in the above definition we join u_1, u_2 with the vertices incident to different edges f_1 and f_2 (then v_1 and v_2 are also joined with the vertices of different edges f_1 and f_2), the resulting cubic graph is called *the crossed connected sum* of G_1 and G_2 and is denoted by $G_1 \sharp G_2$, see Figure 3.2.

It is clear that the operations of double connected sum and crossed connected sums are not determined uniquely and the result $G_1 \star G_2$ depends on the distinguished edges of two graphs.

Let e be an edge of the connected cubic graph G . We shall say that e is *inessential* (subject to the parameter γ) if $\gamma(G) = \gamma(G - e)$. Otherwise e is called *essential*. It is naturally to ask whether the (oriented) genus is additive under taking the operations of connected sum and double connected

FIGURE 3.2. The crossed connected sum of graphs G_1 and G_2

sum of cubic graphs. In general, the answer is negative. For example, we have $\gamma(K_{3,3}) = 1$ while $\gamma(K_{3,3} \star K_{3,3}) = 1 \neq 2$.

Similarly, the genus is not additive subject to the operation of double connected sum of cubic graphs.

The following assertions show that under certain conditions, the (oriented) genus is subadditive or additive with respect to the operations defined above.

Theorem 3.1. *Let G_1 and G_2 be 2-connected cubic graphs of genus k and l , respectively. Let e and f be distinguished edges of G_1 and G_2 , respectively and $G_1 \star G_2$ be the connected sum of G_1 and G_2 . Then*

$$\gamma(G_1) + \gamma(G_2) \geq \gamma(G_1 \star G_2) \geq \gamma(G_1) + \gamma(G_2) - 1.$$

Moreover if e is inessential in G_1 or f is inessential in G_2 , then

$$\gamma(G_1 \star G_2) = k + l.$$

Proof. Let $\varphi_1: G_1 \rightarrow M_1$ be a minimal embedding of the graph G_1 in the surface M_1 with $\gamma(G_1) = \gamma(M_1)$ and $\varphi_2: G_2 \rightarrow M_2$ be a minimal embedding of the graph G_2 in the surface M_2 with $\gamma(G_2) = \gamma(M_2)$. Cut an open disc D_1 in M_1 containing the edge e of G_1 and an open disc D_2 in M_2 containing the edge f of G_2 . Then join the resulting surfaces M'_1 and M'_2 with a tube t , where one connected component of ∂t is identified with ∂D_1 and the other connected component of ∂t is identified with ∂D_2 . The resulting surface is denoted by M . Drawing the bridge edges h_1 and h_2 in the tube t , we obtain an embedding of the graph $G_1 \star G_2$ into the connected surface M . The inequality $\gamma(G_1) + \gamma(G_2) \geq \gamma(G_1 \star G_2)$ now follows.

We continue with proving the second assertion. Denote the two bridge edges of $G_1 \star G_2$ by h_1 and h_2 . Let $\varphi: G_1 \star G_2 \rightarrow M$ be a minimal embedding of the graph $G_1 \star G_2$ in a closed orientable surface M . Then φ is a 2-cell embedding. Consider a regular neighborhood N_1 of the polyhedron $\varphi(G_1) - e$ in M . Then N_1 is a compact 2-manifold with the boundary ∂M . The compact 2-manifold $M_1 = \overline{M \setminus N_1}$ is decomposed into several connected components $S_1, \dots, S_l$. We also have $\partial(N_1) = \partial(M_1)$. The

connected graph $\varphi(G_2 - f)$ is contained in one such connected component, say S_1 . All other connected components of $M \setminus N_1$ must be open 2-disks. We have $\partial(S_1) = m_1 \sqcup m_2 \sqcup \dots \sqcup m_k$ where each m_i is a circle. The 2-manifolds N_1 and S_1 have each boundary component m_i in common.

We claim that each m_i intersects at least one bridge edge h_i of $G_1 \star G_2$. Otherwise we can cut the manifold M along the circle m_j , not intersecting the bridge edges, past the two holes by discs to obtain another closed orientable 2-manifold M' . Note that $\gamma(M') = \gamma(M) - 1$ and the graph $G_1 \star G_2$ has an embedding in M' . But this contradicts to the assumption that the embedding φ of the graph $G_1 \star G_2$ is minimal.

Denote by Π the rotation system on $G_1 \star G_2$ induced by φ . Consider the facial circuits of the rotation system Π that contains bridge edges h_1 and h_2 . We have the following (alternative) possibilities.

1) There are facial circuits c_1 and c_2 of Π such that c_1 contains h_1 (twice) and c_2 contains h_2 (twice). The corresponding closed faces r_1 and r_2 bounded by c_1 and c_2 , respectively, form two handles in M , H_1 and H_2 . Then $\chi(M) \leq 0$. Cutting M along the meridians m_1 and m_2 of H_1 and H_2 and pasting the holes by discs, we shall obtain two disjoint closed orientable surfaces, M_1 and M_2 . This induces embeddings of the graph $G_1 - e$ in the surface M_1 and the graph $G_2 - f$ in the surface M_2 . We have $\gamma(M) = \gamma(M_1) + \gamma(M_2) + 1$. Since $\gamma(G_1 - e) = \gamma(G_1)$ or $\gamma(G_2 - f) = \gamma(G_2)$, the assertion follows.

2) There are two facial circuits c_1 and c_2 of Π each of which contains both the edges h_1 and h_2 . Fix an orientation on M . Let r_1 and r_2 be the faces of the embedding φ bounded by c_1 and c_2 , respectively. The closed faces r_1 and r_2 glued along the edges h_1 and h_2 form a handle. Removing from M the (open) faces r_1, r_2 together with the edges h_1 and h_2 , we shall obtain two disjoint 2-manifolds, M'_1 and M'_2 with boundaries $\partial M'_1$ and $\partial M'_2$, respectively. Elimination of the edges h_1 and h_2 in $G_1 \star G_2$ leads to a surgery of the rotation system Π and induces actually the rotation systems Π_1 and Π_2 on the graphs $G_1 - e$ and $G_2 - f$, respectively. More precisely, instead of the facial circuits c_1 and c_2 in Π we have two new circuits, d_1 and d_2 , respectively, in Π_1 and Π_2 . We thus have $\gamma(G_1 \star G_2) \geq \gamma(G_1 - e) + \gamma(G_2 - f)$. The rotation systems Π_1 and Π_2 generates embeddings of the graphs $G_1 - e$ and $G_2 - f$ in the surfaces M_1 and M_2 , respectively. By drawing the edge e in the face D_1 bounded by the circuit d_1 and the edge f in the disc D_2 bounded by the circuit d_2 , we obtain embeddings of G_1 into M_1 and G_2 into M_2 . Therefore we have $\gamma(M) \geq \gamma(G_1) + \gamma(G_2)$.

3) There is a unique facial circuit c of Π which contains both the edges h_1 and h_2 twice. Now we proceed just as in the case 1). After surgery of the surface M we shall obtain two disjoint surfaces, M_1 and M_2 , such that

$\gamma(M) = \gamma(M_1) + \gamma(M_2) + 1$. Moreover, G_1 has embedding in M_1 and G_2 has embedding in M_2 . Since $\gamma(G_1 - e) = \gamma(G_1)$ or $\gamma(G_2 - f) = \gamma(G_2)$ we have $\gamma(M) \geq \gamma(G_1) + \gamma(G_2)$ completing the proof of the second assertion.

The inequality $\gamma(G_1 \star G_2) \geq \gamma(G_1) + \gamma(G_2) - 1$ follows directly from the proof of the second assertion through the careful analysis of the cases 1)-3). $\square$

Corollary 3.2. *Let G_1 be a 2-connected cubic graphs with the distinguished edge e . Let e' be a distinguished edge of the graph $K_{3,3}$ and $H = G_1 \star K_{3,3}$ be a connected sum of G_1 and $K_{3,3}$ subject to the edges e and e' . If e is inessential in G_1 , then $\gamma(H) = \gamma(G_1) + 1$. $\square$*

Now take in the graph $K_{3,3}$ an edge e and replace it with two parallel edges, e_1 and e_2 . The resulting cubic graph is denoted by K_1 . It is clear that both e_1 and e_2 are inessential in K_1 . Take e_1 as a distinguished edge of K_1 and consider the connected sum $K_2 = K_1 \star K_{3,3}$. By Corollary 3.2, $\gamma(K_2) = 2$. Iterating this process, we obtain a sequence K_l of 2-connected cubic graphs with $\gamma(K_l) = l$. Note that the order of K_l is equal to $8l - 2$.

Corollary 3.3. *If H is a minimal l -genus graph in the class of 2-connected graphs, then $|H| \leq 8l - 2$. $\square$*

Theorem 3.4. *Let G_1 be a 3-connected cubic graph with the pair of distinguished edges e_1 and e_2 and G_2 be a cyclically 4-edge connected cubic graph with the pair of distinguished edges f_1 and f_2 . Assume that $\gamma(G_1 - e_1) = \gamma(G_1)$ or $\gamma(G_1 - e_2) = \gamma(G_1)$ and $\gamma(G_2 - \{f_1, f_2\}) \geq \gamma(G_2) - 1$. Then $G_1 \star G_2$ is a 3-connected graph and $\gamma(G_1 \star G_2) \geq \gamma(G_1) + \gamma(G_2) - 1$.*

Proof. The fact that the graph $G_1 \star G_2$ is 3-connected does not depend on topological properties of graphs G_1 and G_2 and actually follows from the proof of Theorem 3.14 (see below).

Let ψ be an embedding of the graph $G_1 \star G_2$ in a surface M of minimal genus. Consider a subpolyhedron $P = \psi(G_1 - \{e_1, e_2\})$ in M . Let $N(P)$ be a regular neighborhood of P in M . This is a compact submanifold of M (see, for example [11]) and its boundary $\partial(N(P))$ of $N(P)$ consists of k disjoint circles $c_1, \dots, c_k$.

Let S be a complementary submanifold of $N(P)$ in M . It consists of several connected components S_i , $S = \sqcup_i S_i$. Since the graph $G_2 - \{f_1, f_2\}$ is connected, it is contained in one such component, say S_j . Denote by M_2 the closure of submanifold S_j in M . We have obviously $\partial M_2 \subset \partial(N(P))$, so ∂M_2 is the disjoint union of several circles c_i , i.e. $\partial M_2 = c_{i_1} \sqcup \dots \sqcup c_{i_l}$, where $l \leq k$.

As the embedding of $G_1 * G_2$ is genus minimal, all other connected components of $M \setminus N(P)$ must be open 2-disks. Moreover each boundary component c_i of M_2 must intersect at least one bridge edge h_1, h_2, h_3, h_4 . We may suggest without loss of generality that each bridge edge h_m , $m = 1, \dots, 4$, of $G_1 * G_2$ intersects in M a unique circle c_m at one point and each such intersection is transversal (see also the proof of Theorem 3.5). In particular, ∂M_2 consists of at most 4 circles i.e. $l \leq 4$.

Denote by M_1 the submanifold $M \setminus S_j$. Both the submanifolds M_1 and M_2 are connected and by construction we have $\partial M_1 = \partial M_2 = \sqcup_{s=1}^l c_{i_s}$. Moreover the graph $G_1 - \{e_1, e_2\}$ is embedded in M_1 and $G_2 - \{f_1, f_2\}$ is embedded in M_2 . Glue the boundary components of ∂M_1 and ∂M_2 by discs and denote the obtained surfaces by M'_1 and M'_2 , respectively.

Since $G_1 - \{e_1, e_2\} \subseteq M'_1$, we have that $\gamma(M'_1) \geq \gamma(G_1 - \{e_1, e_2\})$. Moreover, from the inequality $\gamma(G_1 - e_1) \leq \gamma(G_1 - \{e_1, e_2\}) + 1$ and assumption $\gamma(G_1 - e_1) = \gamma(G_1)$ we also get that

$$\gamma(M'_1) \geq \gamma(G_1 - \{e_1, e_2\}) \geq \gamma(G_1 - e_1) - 1 = \gamma(G_1) - 1.$$

As $G_2 - \{f_1, f_2\} \subseteq M'_2$, by the assumption $\gamma(G_2 - \{f_1, f_2\}) \geq \gamma(G_2) - 1$, we have

$$\gamma(M'_2) \geq \gamma(G_2 - \{f_1, f_2\}) \geq \gamma(G_2) - 1.$$

We have to show that $\gamma(M) \geq \gamma(G_1) + \gamma(G_2) - 1$. Suppose

$$\gamma(M) \geq \gamma(G_1) + \gamma(G_2) - 2.$$

By the above reasoning, this is possible whenever we have the following $\gamma(M'_1) = \gamma(G_1) - 1$ and $\gamma(M'_2) = \gamma(G_2) - 1$. In other words,

$$\gamma(M) = \gamma(M_1) + \gamma(M_2),$$

so M is obtained from M_1 and M_2 by gluing along one boundary component. Therefore $\partial M_1 = \partial M_2$ is a circle c . In particular, M'_1 is obtained from M_1 by attaching a disc D , so $\gamma(M'_1) = \gamma(M_1)$. It follows that the both ends of e_2 belong to c , so one can extend embedding $G_1 - \{e_1, e_2\} \subset M_1$ to embedding $G_1 - e_1 \subset M'_1 = M_1 \cup D$ by drawing e_2 in the closed 2-cell $\bar{D}$. We thus get embedding of $G_1 - e_1$ into surface M'_1 of genus $\gamma(G_1) - 1$ contradicting to our assumption. $\square$

Note also that an analogue of Theorem 3.4 holds also for crossed connected sum of cubic graphs.

Let G_1 be a 2-connected cubic graph of genus $k > 0$ which has the pair of distinguished non incident edges $\{e_1 = (u_1, v_1), e_2 = (u_2, v_2)\}$ and let G_2 be a connected cubic graph with the pair of distinguished non incident edges $\{f_1 = (u'_1, v'_1), f_2 = (u'_2, v'_2)\}$. Assume that the following two conditions holds:

- (i) at least one of the edges e_1, e_2 in G_1 is inessential;
- (ii) $\gamma(G_2) = 1$ and either $\gamma(G_2 - \{f_1, f_2\}) = 1$, or $\gamma(G_2 - \{f_1, f_2\}) = 0$

and for any plain embedding of $G_2 - \{f_1, f_2\}$ there is no facial circuit c' containing the four vertices u'_1, v'_1, u'_2, v'_2 and the only possibility that the two facial circuits c'_1, c'_2 cover all these vertices is that one of them contains the vertices u'_1, u'_2 and the other one contains the vertices v'_1, v'_2 .

For a moment, let $G_1 \# G_2$ denote the crossed connected sum of cubic graphs G_1 and G_2 in which the vertices of the pair $\{u_1, v_1\}$ are joined to the vertices of the pair $\{u'_1, u'_2\}$ and the vertices of the pair $\{u_2, v_2\}$ to the vertices of the pair $\{v'_1, v'_2\}$.

Theorem 3.5. *Let G_1 and G_2 be cubic graphs that satisfy conditions (i) and (ii). Assume that G_1 is 3-connected and G_2 is cyclically 4-edge connected. Then $G_1 \# G_2$ is 3-connected graph and $\gamma(G_1 \# G_2) = k + 1$.*

Proof. The proof of the first assertion follows from the proof of the first part of Theorem 3.14.

It remains to prove the second assertion. Suppose that

$$\gamma(G_1 \# G_2) \leq k.$$

Let φ be an embedding of $G_1 \# G_2$ into an orientable surface M of genus k , and ψ be the embedding of the subgraph $G_1 - \{e_1, e_2\}$ into M induced by the embedding φ . Let also $N(G_1)$ be an open regular neighborhood of the polyhedron $\psi(G_1 - \{e_1, e_2\})$ in M .

Let s be a connected component of the 2-manifold $M_2 = M \setminus N(G_1)$ containing the image $\varphi(G_2 - \{f_1, f_2\})$. Then s cannot be a disc (i.e. a face of the embedding ψ). Indeed, otherwise the bridge edges of $G_1 \# G_2$ would join the four vertices from $G_2 - \{f_1, f_2\}$ to four vertices of $G_1 - \{e_1, e_2\}$ in a disc. But this is impossible by condition (ii). Therefore s contains tubes (i.e. is a submanifold with nontrivial fundamental group). It follows that $\gamma(G_1 \# G_2) \geq k$.

It can occur that ∂s consists of one connected component, a circle c . Then $M_1 = \overline{M \setminus s}$ is a 2-manifold with the boundary $\partial M_1 = c$. After gluing a disc D to M_1 along the circle c we shall obtain a surface T of genus $k - 1$. In this case we can draw the edge e_1 (or the edge e_2) in the disc D and obtain an embedding of the graph $G_1 - e_1$ into the surface M_1 contradicting with the equality $\gamma(G_1 - e_1) = k$. We thus exclude this possibility.

Suppose now that s is glued to the rest of the surface M along two or more circles c_i . The number of circles cannot be bigger than two, otherwise the genus of M would be greater than k , contradicting to our assumption.

Assume that s has two boundary components, c_1 and c_2 . Then s is a cylinder and $\gamma(\overline{M \setminus s}) = k - 1$. There are two tubes t_1 and t_2 inside s which contain four bridge edges of the graph $G_1 \# G_2$. A tube t_i , $i = 1, 2$, cannot contain three bridge edges h_i , otherwise one circle c'_i would contain three vertices from the set $L = \{u'_1, v'_1, u'_2, v'_2\}$ and the other circle c'_{3-i} contains the remaining vertex, which is impossible by condition (ii).

Therefore the first tube t_1 , bounded by c_1 on one side, contains two bridge edges h_1 and h_2 joining the ends of the edge e_1 to the vertices, say u'_1 and u'_2 , positioned on the facial circuit c'_1 of $G_2 - f_1 - f_2$.

Similarly, the second tube t_2 , bounded by c_2 on one side, contains the remaining bridge edges h_3 and h_4 which join the ends of the edge e_2 to the vertices v'_1 and v'_2 , positioned on the second facial circuit c'_2 of $G_2 - f_1 - f_2$, see Figure 3.3.

In this case we can add the edges $e_1 = (u_1, v_1)$ and $e_2 = (u_2, v_2)$ to the subgraph $G_1 - e_1 - e_2$ and draw them in the 2-manifold $N(G_1)$. It follows that the graph G_1 admits embedding in a surface of genus $k - 1$ contradicting to the condition (i). This completes the proof of the second assertion. $\square$

FIGURE 3.3.

Example 3.6. Consider the cubic graph K obtained from $K_{3,3}$ by doubling an edge e . Instead of e , we have in K two edges e_1 and e_2 , see Figure 3.4. Take the edges e_1 and e_2 to be distinguished in K . Removing e_1 and e_2 from K we shall obtain a subcubic graph K' . Obviously, K' is homeomorphic to the complete graph K_4 so there is a unique embedding ρ of K' in the sphere S^2 . The pairs of vertices $\{u_1, u_2\}$ and $\{v_1, v_2\}$ are positioned on two different faces of ρ and there is no face r of ρ that contains three of these vertices in the boundary. It follows that K satisfies condition (i) (subject to the pair of edges e_1 and e_2). It is also clear that K satisfies the condition (i) as well (subject to the pair of edges e_1 and e_2).

It follows that $\gamma(K\sharp K) = 2$. Note also that $K\sharp K$ is cyclically 4-edge connected cubic graph.

FIGURE 3.4. The cubic graph K

Lemma 3.7. *The genus of cubic graph $CNG\ 3A$ is equal to 2.*

Proof. Cut the graph $CNG\ 3A$ across four edges as shown in Figure 3.5. We have a decomposition of $CNG\ 3A$ into two planar graphs G_1 and G_2 such that G_1 contains four semiedges e_1, e_2, e_3 and e_4 and G_2 contains four semiedges f_1, f_2, f_3 and f_4 .

FIGURE 3.5.

Suppose that the graph $CNG\ 3A$ is toroidal. Let φ denote embedding of this graph in the torus T . Then φ induces embeddings φ_1 and φ_2 of the subgraphs G_1 and G_2 , respectively, in the torus. Let N_1 and N_2 be open regular neighborhoods of the graphs $\varphi_1(G_1)$ and $\varphi_1(G_2)$, respectively, in T . Then G_1 is contained in one connected component t of the 2-manifold $T \setminus N_2$ and G_2 is contained in one connected component s of the 2-manifold $T \setminus N_1$. The component t cannot be a disc since there is no planar embedding of G_1 which contains all semiedges inside the same region r . Similarly the component s is not a disc. Therefore the only possibility to obtain embedding of the graph $CNG\ 3A$ in the torus is as follows. The subgraph

G_1 is embedding into a sphere S_1 with two holes, the subgraph G_1 is embedding into a sphere S_2 with two holes and the spheres S_1 and S_2 are joining by two tubes τ_1 and τ_2 which contain four pairs of glued semiedges: $(e_1, f_1), (e_2, f_2), (e_3, f_3)$ and (e_4, f_4) . By careful inspection all possibilities we can easily check that this is impossible. $\square$

Now starting from the graphs $CNG3A$ and K in Example 3.6, we can inductively construct a sequence of 3-connected cubic graphs H_l of order $8l$. Note that at each inductive step l , there is at least two nonincident inessential edges in H_l . By Lemma 3.7 and Theorem 3.5 we have $\gamma(H_l) = l$.

Corollary 3.8. *If H is minimal l -genus graph in the class of 3-connected cubic graphs, then $|H| \leq 8l$.*

Denote by $\chi'(G)$ the chromatic index of the graph G . A cubic graph G is called *colorable* if $\chi'(G) = 3$, otherwise G is called *uncolorable* (i.e. $\chi'(G) = 4$) or a *weak snark*. A weak snark which is cyclically 4-edge connected and whose girth is at least five is called a *snark*, [15].

The Petersen graph is a simplest example of a snark. Using the operation of dot product, see Figure 3.6, one obtains from any two snarks of orders k and l , respectively, a bigger snark of order $k + l - 2$. Note that the dot product $G_1 \cdot G_2$ of two cubic graphs G_1 and G_2 is defined non uniquely.

FIGURE 3.6. The dot product of two snarks

In [15] the authors consider different powers P^k of the Petersen graph P and study their genus. A k -th power P^k of the Petersen graph P is defined inductively: $P^k = P \cdot P^{k-1}$, where $\cdot$ denote a dot product of the cubic graphs. Since the dot product of two cubic graphs is defined non uniquely, there are several powers P^n of the snark P for each natural number $n \geq 2$.

In [15] the authors construct for each pair (k, n) of natural numbers k and n , where $k \leq n$ and $k, n \geq 1$, the powers P^n such that $\gamma(P^n) = k$. Note that the order of P^n is equal to $8n + 2$. This is an open problem to evaluate the number $ed(P^n)$ of the powers P^n of P such that $\gamma(P^n) = k$.

In the remaining part of this section, we study additivity properties of the parameter ed subject to operations of connected and double connected sum

of graphs within the classes of 2-connected and 3-connected cubic graphs. A simple example shows that this parameter is not additive under the connected sum of cubic graphs. It suffice to consider the graphs $K_{3,3} * K_{3,3}$ and $K_{3,3} \star K_{3,3}$. Indeed we have $ed(K_{3,3}) = 1$ and $ed(K_{3,3} \star K_{3,3}) = 1$. Moreover $ed(K_{3,3} * K_{3,3}) = 1$ for appropriate choice of pairs of the non incident edges in the first and second copies of $K_{3,3}$. However under certain conditions an analogue of additivity property holds also for the parameter ed .

Let G be a cubic graph and e and f are two distinguished edges of G . We shall say that the edge e of G is *inessential* (subject to the characteristic ed) if $ed(G - e) = ed(G)$.

Example 3.9. Let H denote the Heawood graph. Take any two edges e and f in H , subdivide e with the vertex x and f with the vertex y and connect the new vertices with an edge h . Denote the resulting cubic graph by T . We claim that the edge h is inessential in T . To show this note that $ed(T) = 3$. Indeed the equality $ed(T) = 4$ would imply that $cr(T) = 4$. But the only 4-crossing minimal cubic graphs are Pr_8 and MK . As was mentioned in Section 2, the following equalities hold: $ed(Pr_8) = 2$ and $ed(MK) = 3$. Therefore, we have $ed(T) = 3$. Notice that T cannot be the graph MK , since the latter graph is symmetric, so removing any edge from it and ignoring two new vertices of degree 2, we obtain a cubic graph U of degree 14 and with the girth equal to five. But it is well known that H is of girth six. Note however that h is a unique inessential edge in T , since removal any other edge in T leads to a cubic graph of order 14 that is not isomorphic to H , so to a graph L with $ed(L) = 2$.

Example 3.10. Take in the Heawood graph H an edge e and replace it with two parallel edges, e' and e'' . Let H' denote the resulting cubic graph of order 16. Then both e' and e'' are obviously inessential edges of H' .

Theorem 3.11. *Let G_1 and G_2 be two 2-connected cubic graphs with the distinguished edges e in G_1 and f in G_2 , respectively. If $ed(G_1) = k > 0$ and $ed(G_2) = l > 0$, then $ed(G_1 \star G_2) \geq k + l - 1$. Moreover if e is inessential in G_1 and f is inessential in G_2 , then $ed(G_1 \star G_2) = k + l$.*

Proof. Denote the vertices of e in G_1 by u_1 and u_2 and the vertices of f in G_2 by v_1 and v_2 . Put $ed(G_1 \star G_2) = m$. Let $E = \{e_1, \dots, e_m\}$ be the minimal set of edges in $G_1 \star G_2$ such that $G_1 \star G_2 - E$ is planar. By minimality of E , the graph $G_1 \star G_2 - E$ is connected.

Assume that E contains neither $t_1 = (u_1, v_1)$ nor $t_2 = (u_2, v_2)$. Then either there exists a path p_1 joining u_1 to u_2 in $G_1 - \{e, e_1, \dots, e_m\}$ or a path p_2 in $G_2 - \{f, e_1, \dots, e_m\}$ joining v_1 to v_2 .

Suppose that the first possibility occurs. Then p_1 together with the edges t_1 and t_2 form in $G_1 \star G_2$ a path p that joins v_1 to v_2 and do not intersect $(G_2 - f)$, for exception the end vertices v_1 and v_2 . Evidently, p is homeomorphic to the removed edge f in G_2 and replaces actually it. Since $G_2 - \{f, e_1, \dots, e_m\} \cup p$ is planar, it follows that $|E(G_2 - f) \cap E| \geq l$. As $|E(G_1 - e) \cap E| \geq k - 1$ and $|E(G_2 - f) \cap E| \geq l$, the inequality $|E| \geq k + l - 1$ follows. In the second case, we have $|E(G_1 - e) \cap E| \geq k$ and the assertion also follows.

Assume now that E contains one of the edges t_1 or t_2 . Then E contains at least $k - 1$ edges of $G_1 - e$ and $l - 1$ edges of $G_2 - f$, and the first assertion follows.

The second assertion of the theorem follows directly from the definitions of the connected sum of cubic graphs and the minimal edge deletion set. $\square$

Let G be a connected cubic graph with $ed(G) = l$ and $L \subset E(G)$ be an edge deletion subset of G , so the graph $G - L$ is planar. Note that if the subgraph $G - L$ is disconnected, then $|L| \geq l + 1$. Indeed, suppose contrary that $|L| = l$. We can add some edge r from L to the graph $G - L$ and obtain a planar subgraph U of G . But this contradicts to the assumption that $ed(G) = l$.

Let G be a cyclically 4-edge connected cubic graph with $ed(G) = l$. Let also $e_2 = (u_2, u'_2)$ and $f_2 = (v_2, v'_2)$ be a pair of non incident distinguished and inessential edges in G . Put $G' = G - \{e_2, f_2\}$. Consider an l -cut L in G' which decomposes the graph G' into two planar components, say G'_1 and G'_2 . It may occur that in a plane embedding of $G'_1 \sqcup G'_2$ the pair of vertices $\{u_2, v_2\}$ ($\{u_2, v'_2\}$, respectively) are in the same facial cycle of G'_1 and the pair of vertices $\{u'_2, v'_2\}$ ($\{u'_2, v_2\}$, respectively) are in the same facial cycle of G'_2 . Then L is called a *cut separating* $\{u_2, v_2\}$ from $\{u'_2, v'_2\}$ ($\{u_2, v'_2\}$ from $\{u'_2, v_2\}$, respectively). We shall say that G has the *property P* (subject to the pair of edges e_2 and f_2) if no such separating l -cut L exists in G' .

Example 3.12. Let H' be a cubic graph as in Example 3.10 and

$$\{e' = (u, u'), \quad e'' = (v, v')\}$$

be a pair of distinguished edges in H' . We assert that H' has the property P subject to the pair of edges $\{e', e''\}$. Indeed, first note that H' is cyclically 4-edge connected and $ed(H') = 3$. Remove the edges e' and e'' from H' and denote the resulting graph G' , see Figure 3.7.

The graph G' is also cyclically 4-edge connected and $ed(G') = 2$. Indeed, suppose that G does not have the property P . Let L be a 3-cut of G' that separates $\{u, v\}$ from $\{u', v'\}$ and G_1 and G_2 the corresponding planar components of $G' - L$ such that $u, v \in G_1$ and $u', v' \in G_2$. The only

FIGURE 3.7.

possibility for such a cut is that L contains both the edges f_1, f_2 (so G_1 is simply K_2) and some other edge f of G' . Consider the component K of $G' - \{f_1, f_2\}$ that contains the pair of vertices $\{u', v'\}$. K is of order 14 and contains four vertices of degree two, see Figure 3.8.

FIGURE 3.8.

We can ignore the vertices of degree two in K and consider the corresponding cubic graph K' of degree 10 which is homeomorphic to K . It is easy to see that $g(K') = 5$. It is known that the only cubic graph of degree 10 and of girth 5 is the Petersen graph P . However $ed(P) = 2$, so removal any edge u from it does not lead to a planar graph. It follows that removal any edge f from K does not lead to a planar graph. This contradicts to our assumption that $G_2 = K - f$ is planar. It is rather obvious fact that there is no 3-cut in G' separating $\{u, v\}$ from $\{u', v'\}$. Therefore H' has the property P subject to the pair of edges $\{e', e''\}$.

Example 3.13. Let MK denote the Möbius-Kantor graph. It is well known that MK is a symmetric graph. Take in the graph MK any edge e remove it from MK . Forgetting two vertices of degree 2 in the resulting graph, we obtain a cubic graph U of order 14 with $g(U) = 5$. Then U is not isomorphic to the Heawood graph, so by the results of Section 2 we have $ed(U) = 2$. It follows that each edge of MK is essential. Now replace the edge e in MK with two parallel edges $\{e' = (u, u'), e'' = (v, v')\}$. Denote the resulting graph by W . It is easy to see that both e' and e'' are

inessential edges of W . Let e' and e'' be the distinguished edges of W . By the construction, W is cyclically 4-edge connected and $ed(W) = 3$.

Moreover, we assert that W has the property P subject to the pair of edges $\{e', e''\}$. Indeed, suppose the contrary. Remove the edges e' and e'' from W and denote the resulting graph W' , see Figure 3.9. Since W' differs from U by subdivision only, it is cyclically 4-edge connected and we have $ed(W') = 2$.

FIGURE 3.9.

Let M be a 3-cut of W' that separates $\{u, v\}$ from $\{u', v'\}$ and W_1 and W_2 the corresponding planar components of $W' - M$ where $u, v \in W_1$ and $u', v' \in W_2$. The only possibility for such a cut is that M contains both the edges f_1, f_2 (so W_1 is simply K_2) and some other edge f of W' . Consider the component R of $W' - \{f_1, f_2\}$ that contains the pair of vertices $\{u', v'\}$ (up to homeomorphism of graphs, to obtain R from MK we simply remove from it a star of vertex i.e. the graph $K_{1,3}$). As before, we can forget the vertices u' and v' of degree two in R and consider the corresponding cubic graph R' of degree 12 that is homeomorphic to R . It is easy to see that $g(R') = 5$, see Figure 3.10. It is not difficult to check that $ed(R') = 2$. It follows that removing an edge f from R does not lead to a planar graph. This contradicts to our assumption that $W_2 = R - f$ is planar. It is rather an obvious fact that there is no 3-cut in W' separating $\{u, v\}$ from $\{u', v'\}$. Therefore W has the property P subject to the pair of edges $\{e', e''\}$.

FIGURE 3.10.

Theorem 3.14. *Let G_1 be a 3-connected cubic graph with $ed(G_1) = k > 0$ and G_2 a cyclically 4-edge connected cubic graph with $ed(G_2) = l > 0$. Let $\{e_1, f_1\}$ be a pair of distinguished non incident edges in G_1 and $\{e_2, f_2\}$ a pair of non incident distinguished edges in G_2 . Assume that in both the pairs each edge is inessential and G_1 has the property P with respect to the pair $\{e_1, f_1\}$ and G_2 has the property P with respect to the pair $\{e_2, f_2\}$. Then $G_1 * G_2$ is a 3-connected cubic graph and $ed(G_1 * G_2) \geq k + l$.*

Proof. Let $e_1 = (u_1, u'_1), f_1 = (v_1, v'_1), e_2 = (u_2, u'_2)$ and $f_2 = (v_2, v'_2)$. The bridge edges in the graph $H = G_1 * G_2$ are the following:

$$h_1 = (u_1, u_2), \quad h_2 = (u'_1, u'_2), \quad h_3 = (v_1, v_2), \quad h_4 = (v'_1, v'_2).$$

The subgraph of H formed by the bridge edges h_1, h_2, h_3, h_4 is denoted by B . Put $G'_1 = G_1 - \{e_1, f_1\}$ and $G'_2 = G_2 - \{e_2, f_2\}$.

Note that if G is cyclically 4-edge connected cubic graph, then any 3-edge cut of G is of kind $K_{1,3}$ (see, for example, [22]). We claim that the graph H is 3-connected. Suppose contrary that $\lambda(H) = 2$. Let $A = \{a_1, a_2\}$ be a cut of H consisting of two edges a_1 and a_2 . Depending on the positions of edges a_1 and a_2 in H , the following situations can occur.

1) $a_1 \in E(G'_1)$ and $a_2 \in E(G'_2)$. Under this assumption we have

$$\lambda(G'_1) = \lambda(G'_2) = 1,$$

so the graph H is decomposed into two components by removing the edges a_1 and a_2 as shown in the Figure 3.11. It follows that $E_1 = \{e_1, f_1, a_1\}$ is a 3-cut of the graph G_1 and $E_2 = \{e_2, f_2, a_2\}$ is a 3-cut of the graph G_2 . However since G_2 is cyclically 4-edge connected, this is impossible because the graph formed by the set of edges E_2 is not isomorphic to $K_{1,3}$.

FIGURE 3.11.

2) Both a_1 and a_2 are the edges of G'_1 . In this case, there are paths p_1 and p_2 in $H - E(G'_1)$ that join the vertices u_1, u'_1 and v_1, v'_1 , respectively. This means that U decomposes G'_1 into two subgraphs, say T_1 and T_2 , so that one such T_i contains all the vertices u_1, u'_1, v_1, v'_1 . This would imply that $\lambda(G_1) = 2$ contradicting to our assumption.

3) Both a_1 and a_2 are the edges of G'_2 . This case can be handled just in the same way as the case 2).

Finally, it is obviously that H does not possess a 2-cut which contains at least one bridge edge. Therefore H is 3-connected. It remains to show that $ed(G_1 * G_2) \geq k + l$.

Let $R = \{r_1, \dots, r_s\}$ be a minimal edge deletion set in H . Denote by q the number $|R \cap \{h_1, h_2, h_3, h_4\}|$. The planar graph $H - R$ is connected. The proof of the inequality is actually reduced to analyzing the following three cases:

Case 1) $q \geq 2$. By assumption, we have

$$ed(G'_1) \geq k - 1 \quad \text{and} \quad ed(G'_2) \geq l - 1.$$

Since the graphs G'_1 and G'_2 are planar, it follows that

$$|E(G'_1) \cap R| \geq ed(G'_1) \geq k - 1, \quad |E(G'_2) \cap R| \geq ed(G'_2) \geq l - 1.$$

As the sets $E(G'_1) \cap R$ and $E(G'_2) \cap R$ are disjoint, we have

$$|R| \geq k - 1 + l - 1 + q \geq k + l,$$

so $ed(H) \geq k + l$.

Case 2) $q = 0$, i.e. R does not contain any bridge edge h_i . Consider a planar drawing g of the connected graph $H - R$. Let g_1 and g_2 be the planar embeddings of the subgraphs $G'_1 - R$ and $G'_2 - R$, respectively, induced by g . To prove the assertion in this case we have to inspect the following three subcases.

(i) *Both the subgraphs $G'_1 - R$ and $G'_2 - R$ are connected.* Since $G'_1 - R$ is connected, the plane subgraph $D_2 = (G'_2 \cup B) - R$ is contained in a face μ of the plane embedding g_1 of the graph $G'_1 - R$. This means that the vertices u_1, u'_1, v_1, v'_1 of $G'_1 - R$ are situated on the same facial circuit c , the circuit that bounds the face μ . We can draw the edge e_1 (or the edge f_1) in the face μ and obtain a planar embedding of the subgraph $G_1 - (f_1 \cup R)$, see Figure 3.12. Since the edge f_1 is inessential in G_1 , we have $|E(G'_1) \cap R| \geq k$. In the same way we can prove that $|E(G'_2) \cap R| \geq l$. It follows that $|R| \geq k + l$.

(ii) *Both the subgraphs $G'_1 - R$ and $G'_2 - R$ are disconnected.* Then $|E(G'_2) \cap R| \geq l$ and $|E(G'_1) \cap R| \geq k$, so $|R| = k + l$;

(iii) *One of the subgraphs $G'_1 - R$ and $G'_2 - R$ is connected and the other is disconnected.* Suppose for instance that $G'_1 - R$ is connected and $G'_2 - R$ is disconnected. Then $G'_2 - R$ consists of two connected components, say U_1 and U_2 . Since $G'_2 - R$ is disconnected, we have $|E(G'_2) \cap R| = l$. If $|E(G'_1) \cap R| = k$ we have $|R| = k + l$. Suppose that $|E(G'_1) \cap R| = k - 1$. Then in any plane embedding of $W = G'_1 - R$, the pair of vertices u_1 and u'_1 and the pair of vertices v_1, v'_1 cannot be neighboring (positioned on the

FIGURE 3.12.

same face). For the given plane subgraphs $G'_2 - R$ and $G'_1 - R$, the only possibility to draw the graph H in the plane is to take a face c_1 of $G'_1 - R$ containing u_1 and v_1 , a face c_2 containing u'_1 and v'_1 (if such exist), insert a connected component U_i into the face c_1 , the second connected component U_{3-i} into the face c_2 , then connect the vertices u_1 with u_2 and the vertices v_1 with v_2 with bridge edges h_1 and h_3 inside c_1 , and the vertices u'_1 with u'_2 and the vertices v'_1 with v'_2 inside the face c_2 with bridge edges h_2 and h_4 , see Figure 3.13. This is possible only if u_2 and v_2 are vertices of the same face of the plane graph U_i and u'_2 and v'_2 are vertices of the same face of the plane graph U_{3-i} . However the latter is excluded by property P . Therefore $|E(G'_1) \cap R| = k$ and $|R| = k + l$.

FIGURE 3.13.

Case 3) $q = 1$. If $|E(G'_2) \cap R| \geq l$ or $|E(G'_1) \cap R| \geq k$, the assertion follows. Suppose that $|E(G'_2) \cap R| = l - 1$ and $|E(G'_1) \cap R| = k - 1$. By the same arguments as before we conclude that in this case both the graphs $G'_1 - R$ and $G'_2 - R$ are connected. Let g be a planar embedding of the connected graph $H - R$. The embedding g induces planar embeddings g_1 and g_2 of the subgraphs $G'_1 - R$ and $G'_2 - R$, respectively. Since $G'_1 - R$ is connected, the plane subgraph $D_2 = (G'_2 \cup B) - R$ is contained in a face γ of the plane graph $G'_1 - R$. This means that three vertices of $G'_1 - R$ from the set $\{u_1, u'_1, v_1, v'_1\}$ are situated in the same facial circuit c , the circuit

that bounds the face γ . Therefore we can draw the edge e_1 or the edge f_1 in the face γ which gives in planar embedding of the subgraph $G_1 - (f_1 \cup R)$ ($G_1 - (e_1 \cup R)$, respectively). Since the edges e_1 and f_1 are inessential in G_1 we have $|E(G'_1) \cap R| \geq k$, contradicting to our assumption.

We conclude that in any case $|R| \geq k + l$. We thus have proved that $ed(H) \geq k + l$. $\square$

Theorem 3.14 can be used in constructing 3-connected and even cyclically 4-edge connected cubic graphs G of order $16n$ such that $ed(G) \geq 3n$ for any natural number $n > 0$. We can start from the graph H' as in Example 3.10. Let e' and e'' be two parallel edges in H' obtained by doubling an edge e in the Heawood graph H . As was noted before, H is cyclically 4-connected and $\zeta(H') = 4$. Taking two copies of H' , the graphs G' and G_1 , and applying to them the operation of double connected sum (just as it was described in Theorem 3.14), we shall obtain a 3-connected cubic graph H_2 of order 32. By Theorem 3.14, since H' has the property P , we get that $ed(H_2) = 6$. It is not difficult to check that $\zeta(H_2) = 4$. The thorough analysis of the proof of Theorem 3.14 shows that each bridge edge h_i in H_2 is also inessential.

Now we can take three copies of the cubic graph H' , the graphs U_1 , U_2 and U_3 with the pairs of distinguished edges e_1, f_1 , and e_2, f_2 and e_3, f_3 , respectively, remove all them and join the resulting graphs G_1, G_2 and G_3 by six bridge edges h_i as shown in Figure 3.14. The resulting cubic graph H_3 is 3-connected and we have $ed(H_3) = 9$. The proof of this assertion actually follows from the proof of Theorem 3.14 and uses in an essential way the facts that H' is cyclically 4-edge connected and H has the property P . We omit here the details.

FIGURE 3.14. The cubic graph H_3

Iterating the process of joining the several copies of the graph H' in a cycle in the way as before, we obtain a sequence of cyclically 4-edge connected cubic graphs H_n of order $16n$ with $ed(H_n) = 3n$.

Corollary 3.15. *If H is an $3l$ -edge deletion minimal graph in the class of 3-connected cubic graphs, then $|H| \leq 16l$.*

Question. We provided some upper bounds for the order of minimal edge deletion (cubic) graphs and minimal genus (cubic) graphs. What about nontrivial lower bounds for these graph parameters?

Acknowledgements. The author would like to thank the referee for useful remarks and comments and for correcting some mistakes in the previous version of the manuscript.

REFERENCES

- [1] Joseph Battle, Frank Harary, Yukihiro Kodama, J. W. T. Youngs. Additivity of the genus of a graph. *Bull. Amer. Math. Soc.*, 68:565–568, 1962.
- [2] Daniel Bienstock, Nathaniel Dean. Bounds for rectilinear crossing numbers. *J. Graph Theory*, 17(3):333–348, 1993.
- [3] Gruia Călinescu, Cristina G Fernandes, Ulrich Finkler, Howard Karloff. A better approximation algorithm for finding planar subgraphs. *J. Algorithms*, 27(2):269–302, May 1998.
- [4] Luerbio Faria, Celina M. Herrera de Figueiredo, Sylvain Gravier, Candido F. X. de Mendonça, Jorge Stolfi. On maximum planar induced subgraphs. *Discrete Appl. Math.*, 154(13):1774–1782, 2006.
- [5] Luerbio Faria, Celina M. Herrera de Figueiredo, Candido F. X. Mendonça. On the complexity of the approximation of nonplanarity parameters for cubic graphs. *Discrete Appl. Math.*, 141(1-3):119–134, 2004.
- [6] Jean-Luc Fouquet, Henri Thuillier. On some conjectures on cubic 3-connected graphs. *Discrete Math.*, 80(1):41–57, 1990.
- [7] M. R. Garey, D. S. Johnson. Crossing number is NP-complete. *SIAM J. Algebraic Discrete Methods*, 4(3):312–316, 1983.
- [8] Jonathan L. Gross. Embeddings of cubic Halin graphs: genus distributions. *Ars Math. Contemp.*, 6(1):37–56, 2013.
- [9] Jonathan L. Gross, Thomas W. Tucker. *Topological graph theory*. Dover Publications, Inc., Mineola, NY, 2012.
- [10] R. K. Guy. A combinatorial problem. *Nabla (Bulletin of the Malayan Mathematical Society)*, 7:68–72, 1960.
- [11] J. F. P. Hudson. *Piecewise linear topology*. University of Chicago Lecture Notes prepared with the assistance of J. L. Shaneson and J. Lees. W. A. Benjamin, Inc., New York-Amsterdam, 1969.
- [12] Ed Pegg Jr, G. Exoo. Crossing number graphs. *The Mathematica Journal*, 11(2):161–170, 2009.
- [13] Gary L. Miller. An additivity theorem for the genus of a graph. *J. Combin. Theory Ser. B*, 43(1):25–47, 1987.
- [14] Bojan Mohar, Carsten Thomassen. *Graphs on surfaces*. Johns Hopkins Studies in the Mathematical Sciences. Johns Hopkins University Press, Baltimore, MD, 2001.
- [15] Bojan Mohar, Andrej Vodopivec. The genus of Petersen powers. *J. Graph Theory*, 67(1):1–8, 2011.
- [16] Roman Nedela, Martin Škoviera. Atoms of cyclic connectivity in cubic graphs. *Math. Slovaca*, 45(5):481–499, 1995.
- [17] Shengjun Pan, R. Bruce Richter. The crossing number of K_{11} is 100. *J. Graph Theory*, 56(2):128–134, 2007.
- [18] L. Plachta. On minimal genus cubic graphs. *in preparation*.

- [19] Gerhard Ringel. *Map color theorem*. Springer-Verlag, New York-Heidelberg, 1974. Die Grundlehren der mathematischen Wissenschaften, Band 209.
- [20] László A. Székely. A successful concept for measuring non-planarity of graphs: the crossing number. *Discrete Math.*, 276(1-3):331–352, 2004. 6th International Conference on Graph Theory.
- [21] Carsten Thomassen. The graph genus problem is NP-complete. *J. Algorithms*, 10(4):568–576, 1989.
- [22] Nicholas C. Wormald. Classifying k -connected cubic graphs. In *Combinatorial mathematics, VI (Proc. Sixth Austral. Conf., Univ. New England, Armidale, 1978)*, volume 748 of *Lecture Notes in Math.*, 199–206. Springer, Berlin, 1979.

Received: September 14, 2017, accepted: June 20, 2018.

Leonid Plachta

AGH UNIVERSITY OF SCIENCE AND TECHNOLOGY (AL. MICKIEWICZA 30, 30-059 CRACOW, POLAND), & INSTITUTE OF APPLIED PROBLEMS OF MECHANICS AND MATHEMATICS OF NAS OF UKRAINE (3-B NAUKOVA ST., 79060, LVIV, UKRAINE)

Email: dept25@gmail.com